

JS006080553A

United States Patent [19]

Sogabe et al.

[11] Patent Number: 6,080,553 [45] Date of Patent: *Jun. 27, 2000

[54]	CREATINE AMIDINOHYDROLASE, PRODUCTION THEREOF AND USE THEREOF			
[75]	Inventors:	Atsushi Sogabe; Takashi Hattori; Yoshiaki Nishiya; Yoshihisa Kawamura, all of Tsuruga, Japan		
[73]	Assignee:	Toyo Boseki Kabushiki Kaisha, Osaka, Japan		
[*]	Notice:	This patent issued on a continued prosecution application filed under 37 CFR 1.53(d), and is subject to the twenty year patent term provisions of 35 U.S.C. 154(a)(2).		
[21]	Appl. No.:	: 08/799,897		
[22]	Filed:	Feb. 13, 1997		
[30]	Forei	gn Application Priority Data		
Feb.	13, 1996	[JP] Japan 8-025435		
[51]	Int. Cl. ⁷	C12Q 1/34; C12N 9/78; C12N 1/20; C12N 1/00		
[52]				
[58]	Field of S	Search		
[56]		References Cited		
U.S. PATENT DOCUMENTS				

3,806,420 4/1974 Holz et al. 195/66

3,907,644	9/1975	Mollering et al.	195/99
5.451.520	9/1995	Furukawa et al.	435/227

FOREIGN PATENT DOCUMENTS

62 091182 4/1987 Japan . 07 265074 10/1995 Japan .

Primary Examiner—Elizabeth Slobodyansky Attorney, Agent, or Firm—Leydig, Voit & Mayer, Ltd.

ABSTRACT

A creatine amidinohydrolase having the following physicochemical properties:

Action: catalyzing the following reaction; creatine+ $H_2O \rightarrow sarcosine+urea$

Optimum temperature: about 40-50° C.

Optimum pH: pH about 8.0-9.0

Heat stability: not more than about 50° C. (pH 7.5, 30 min)

Km value for creatine in a coupling assay using a sarcosine oxidase and a peroxidase: about 3.5-10.0 mM

Molecular weight: about 43,000 (SDS-PAGE)

Isoelectric point: about 3.5,

a method for producing said enzyme, comprising culture of microorganism producing said enzyme, a method for the determination of creatine or creatinine in a sample using said enzyme, and a reagent therefor.

23 Claims, 2 Drawing Sheets